arXiv:1211.4999v1 [math.PR] 21 Nov 2012

SUBSIGNATURES OF SYSTEMS

JEAN-LUC MARICHAL

Abstract. We introduce the concept of subsignature for semicoherent systems as a class of indexes that range from the system signature to the Barlow-Proschan importance index. Specifically, given a nonempty subset M of the set of components of a system, we define the M-signature of the system as the $|M|$-tuple whose kth coordinate is the probability that the kth failure among the components in M causes the system to fail. We give various explicit linear expressions for this probability in terms of the structure function and the distribution of the component lifetimes. We also examine the case of exchangeable lifetimes and the special case when the lifetime are i.i.d. and M is a modular set.

1. Introduction

Let $\mathcal{S} = (C, \phi, F)$ be an n-component system, where $C = \{1, \ldots, n\}$ denotes the set of components, ϕ denotes the associated structure function $\phi\colon \{0,1\}^n \to \{0,1\}$ (which expresses the state of the system in terms of the states of its components), and F denotes the joint c.d.f. of the component lifetimes $T_1, \ldots, T_n$. We assume that the system is *semicoherent*, i.e., the structure function ϕ is nondecreasing in each variable and satisfies the conditions $\phi(0, \ldots, 0) = 0$ and $\phi(1, \ldots, 1) = 1$. We also assume that the c.d.f. F has no ties, that is, $\Pr(T_i = T_j) = 0$ for all distinct $i, j \in C$.

The *signature* of the system, a concept introduced first in 1985 by Samaniego [9] for systems whose components have continuous and i.i.d. lifetimes and then recently extended to non-i.i.d. lifetimes (see [5] and the references therein), is defined as the n-tuple $\mathbf{p} = (p_1, \ldots, p_n)$, where p_k is the probability that the kth component failure causes the system to fail. That is,

$$p_k \;=\; \Pr(T_S = T_{k:n}), \qquad k \in \{1, \ldots, n\},$$

where T_S denotes the system lifetime and $T_{k:n}$ denotes the kth smallest lifetime, i.e., the kth order statistic obtained by rearranging the variables $T_1, \ldots, T_n$ in ascending order of magnitude.

In the i.i.d. case (and even when the component lifetimes are exchangeable) the system signature $\mathbf{p}$ is independent of the c.d.f. F. It is then called the *structural signature* and denoted by $\mathbf{s} = (s_1, \ldots, s_n)$, where $s_k = \Pr(T_S = T_{k:n})$. In fact, Boland

---

Date: November 21, 2012.

2010 *Mathematics Subject Classification.* 62N05, 90B25, 94C10.

Key words and phrases. Semicoherent system, dependent lifetimes, Barlow-Proschan index, system signature, system subsignature.

[3] showed that s_k can be explicitly written in the form

$$s_k = \sum_{\substack{A\subseteq C\\ |A|=n-k+1}} \frac{1}{\binom{n}{|A|}}\,\phi(A) - \sum_{\substack{A\subseteq C\\ |A|=n-k}} \frac{1}{\binom{n}{|A|}}\,\phi(A)\,. \tag{1}$$

Here and throughout we identify Boolean vectors $\mathbf{x}\in\{0,1\}^n$ and subsets $A\subseteq C$ by setting $x_i=1$ if and only if $i\in A$. We thus use the same symbol to denote both a function $f\colon\{0,1\}^n\to\mathbb{R}$ and its corresponding set function $f\colon 2^C\to\mathbb{R}$, interchangeably. For instance, we have $\phi(C)=\phi(1,\ldots,1)$.

Marichal and Mathonet [5] extended formula (1) to the non-i.i.d. case into

$$p_k = \sum_{\substack{A\subseteq C\\ |A|=n-k+1}} q(A)\,\phi(A) - \sum_{\substack{A\subseteq C\\ |A|=n-k}} q(A)\,\phi(A)\,, \tag{2}$$

where the function $q\colon 2^C\to[0,1]$, called the *relative quality function* associated with F, is defined by

$$q(A) = \Pr\Big(\max_{i\in C\smallsetminus A} T_i < \min_{i\in A} T_i\Big).$$

That is, $q(A)$ is the probability that the best $|A|$ components are precisely those in A.

Since its introduction the concept of signature proved to be a very useful tool in the analysis of semicoherent systems, especially for the comparison of different system designs and the computation of the system reliability (see, e.g., [10] for the i.i.d. case and [5, 7] for the non-i.i.d. case).

The *Barlow-Proschan importance index* of the system, another useful concept introduced first in 1975 by Barlow and Proschan [1] for systems whose components have continuous and independent lifetimes and then extended to the general dependent case in [4, 6], is defined as the n-tuple $\mathbf{I}_{\mathrm{BP}}$ whose jth coordinate is the probability that the failure of component j causes the system to fail, that is,

$$I_{\mathrm{BP}}^{(j)} = \Pr(T_S = T_j)\,.$$

Just as for the signature, in the i.i.d. case (and even when the component lifetimes are exchangeable) the index $\mathbf{I}_{\mathrm{BP}}$ is also independent of the c.d.f. F. It is then called the *structural importance index* and denoted $\mathbf{b}=(b_1,\ldots,b_n)$, where $b_j=\Pr(T_S=T_j)$. An explicit expression for b_j in terms of the structure function values is given by

$$b_j = \sum_{A\subseteq C\smallsetminus\{j\}} \frac{1}{n\binom{n-1}{|A|}}\,\Delta_j\phi(A)\,, \tag{3}$$

where $\Delta_j\phi(A)=\phi(A\cup\{j\})-\phi(A\smallsetminus\{j\})$. Marichal and Mathonet [6] extended this formula to the non-i.i.d. case into

$$I_{\mathrm{BP}}^{(j)} = \sum_{A\subseteq C\smallsetminus\{j\}} q_j(A)\,\Delta_j\phi(A)\,, \tag{4}$$

where the function $q_j\colon 2^{C\smallsetminus\{j\}}\to[0,1]$ is defined by

$$q_j(A) = \Pr\Big(\max_{i\in C\smallsetminus A} T_i = T_j < \min_{i\in A} T_i\Big). \tag{5}$$

That is, $q_j(A)$ is the probability that the components that are better than component j are precisely those in A.

For instance, when $n = 4$ we have

$$\begin{aligned} q_2(\{1,3\}) &= \Pr(T_4 < T_2 < \min\{T_1,T_3\}) \\ &= \Pr(T_4 < T_2 < T_1 < T_3) + \Pr(T_4 < T_2 < T_3 < T_1)\,. \end{aligned}$$

More generally, (5) can be rewritten as

$$q_j(A) \;=\; \sum_{\substack{\sigma\in\mathfrak{S}_n\,:\,\{\sigma(n-|A|+1),\ldots,\sigma(n)\}=A\\ \sigma(n-|A|)=j}} \Pr(T_{\sigma(1)} < \cdots < T_{\sigma(n)})\,, \tag{6}$$

where $\mathfrak{S}_n$ denotes the set of permutations on C.

These important concepts of signature and Barlow-Proschan index motivate the introduction of the following more general concept. Let M be a nonempty subset of the set C of components and let $m = |M|$. We define the *M-signature* of the system as the m-tuple $\mathbf{p}^M = (p_1^M,\ldots,p_m^M)$, where p_k^M is the probability that the kth failure among the components in M causes the system to fail. That is,

$$p_k^M \;=\; \Pr(T_S = T_{k:M}), \qquad k\in\{1,\ldots,m\}\,,$$

where $T_{k:M}$ denotes the kth smallest lifetime of the components in M, i.e., the kth order statistic obtained by rearranging the variables T_i ($i\in M$) in ascending order of magnitude. A *subsignature* of the system is an M-signature for some $M\subseteq C$. We naturally denote the probability p_k^M by s_k^M whenever the component lifetimes are exchangeable.

Clearly, when $M = C$ the M-signature reduces to the signature $\mathbf{p}$, which shows that the signature is a particular subsignature. At the opposite, when M is a singleton $\{j\}$ the M-signature reduces to the 1-tuple $\mathbf{p}^{\{j\}} = (p_1^{\{j\}})$, where $p_1^{\{j\}} = \Pr(T_S = T_j)$ is the jth coordinate of the Barlow-Proschan index. Thus, the subsignatures define a class of 2^n-1 indexes that range from the standard signature (when $M = C$) to the Barlow-Proschan index (when M consists of a single component).

Remark 1. The concept of M-signature is particularly relevant for instance when the system has a huge number of components and M is a small subset of potentially unreliable components.

Interestingly, we have the following link between the subsignatures and the Barlow-Proschan index. For every nonempty subset $M\subseteq C$, we have

$$\sum_{k=1}^m p_k^M \;=\; \Pr(T_S = T_j \text{ for some } j\in M) \;=\; \sum_{j\in M} I_{\mathrm{BP}}^{(j)}\,.$$

If this probability is not zero, then we can express the *normalized M-signature* $p_k^M/\sum_{\ell=1}^m p_\ell^M$ as the conditional probability

$$\frac{p_k^M}{\sum_{\ell=1}^m p_\ell^M} \;=\; \Pr(T_S = T_{k:M} \mid T_S = T_j \text{ for some } j\in M)\,.$$

Example 1. Consider the six-component system indicated in Figure 1, where the components have i.i.d. lifetimes. The corresponding structure function is given by

$$\phi(x_1,\ldots,x_6) \;=\; (x_1 \amalg x_2)\, x_3\, x_4\, (x_5 \amalg x_6)\,,$$

where $\amalg$ is the coproduct operation defined by $x \amalg y = 1-(1-x)(1-y)$.

Table 1 gives the normalized subsignatures for some subsets (those for the remaining subsets can be derived by symmetry). These normalized subsignatures illustrate the following fact: The more concentrated the distribution of s^M is on

M	$s^M/\sum_{k=1}^m s_k^M$
$\{1,2\}$	(0.000, 1.000)
$\{1,3\}$	(0.846, 0.154)
$\{3,4\}$	(1.000, 0.000)
$\{1,2,3\}$	(0.467, 0.533, 0.000)
$\{1,3,4\}$	(0.833, 0.167, 0.000)
$\{1,2,5\}$	(0.167, 0.666, 0.167)
$\{1,2,3,4\}$	(0.538, 0.462, 0.000, 0.000)
$\{1,2,5,6\}$	(0.000, 0.500, 0.500, 0.000)
$\{1,2,4,5\}$	(0.412, 0.471, 0.118, 0.000)
$\{1,3,4,5\}$	(0.692, 0.269, 0.038, 0.000)
$\{1,2,3,4,5\}$	(0.464, 0.429, 0.107, 0.000, 0.000)
$\{1,2,3,5,6\}$	(0.263, 0.421, 0.316, 0.000, 0.000)
$\{1,2,3,4,5,6\}$	(0.333, 0.400, 0.267, 0.000, 0.000, 0.000)

TABLE 1. Normalized subsignatures for representative subsets

the left, the more M is "critical" in the sense that each of its components makes the system vulnerable and potentially unreliable. □



FIGURE 1. A six-component system

In this paper we give explicit linear expressions for M-signatures, including a simultaneous generalization of formulas (2) and (4) as well as a linear expression in terms of the Möbius transform of the structure function (Section 2). We also investigate the special case when the component lifetimes are exchangeable (Section 3). Finally, we examine the case when the lifetimes are i.i.d. and M is a modular set (Section 4).

2. EXPLICIT EXPRESSIONS

In this section we provide various explicit expressions for the probability $p_k^M = \Pr(T_S = T_{k:M})$. We start with a linear expression in terms of the functions q_j and $\Delta_j\phi$, thus generalizing formula (4).

Theorem 2. *For every nonempty set $M \subseteq C$ and every $k \in \{1,\ldots,m\}$, we have*

$$p_k^M = \sum_{\substack{A\subseteq C\\|M\setminus A|=k}} \sum_{j\in M\setminus A} q_j(A)\,\Delta_j\,\phi(A) = \sum_{j\in M} \sum_{\substack{A\subseteq C\setminus\{j\}\\|M\setminus A|=k}} q_j(A)\,\Delta_j\,\phi(A)\,. \tag{7}$$

Proof. Fix $\sigma \in \mathfrak{S}_n$, set $j \in C$ such that $T_j = T_{k:M}$, and let $j' = \sigma^{-1}(j)$. Then we have

$$\Pr(T_S = T_j \mid T_{\sigma(1)} < \cdots < T_{\sigma(n)}) = \phi\big(\{\sigma(j'),\ldots,\sigma(n)\}\big) - \phi\big(\{\sigma(j'+1),\ldots,\sigma(n)\}\big)\,.$$

Indeed, the left-hand expression of this equation takes its values in $\{0,1\}$ and is exactly 1 if and only if $\{\sigma(1),\ldots,\sigma(j'-1)\}$ is not a cut set and $\{\sigma(1),\ldots,\sigma(j')\}$ is a cut set. (Recall that a subset $K \subseteq C$ of components is a *cut set* for the function ϕ if $\phi(C \setminus K) = 0$.) That is,

$$\phi\big(\{\sigma(j'), \ldots, \sigma(n)\}\big) = 1 \quad \text{and} \quad \phi\big(\{\sigma(j'+1), \ldots, \sigma(n)\}\big) = 0.$$

Since j' depends on σ, we denote it by j'_σ. By using the law of total probability, we then obtain

$$\Pr(T_S = T_{k:M}) \;=\; \sum_{\sigma \in \mathfrak{S}_n} \Big(\phi\big(\{\sigma(j'_\sigma), \ldots, \sigma(n)\}\big) - \phi\big(\{\sigma(j'_\sigma+1), \ldots, \sigma(n)\}\big)\Big) \Pr(T_{\sigma(1)} < \cdots < T_{\sigma(n)})\,.$$

Grouping the terms for which $\{\sigma(j'_\sigma), \ldots, \sigma(n)\}$ is a union $B_1 \cup B_2 \cup \{i\}$ of fixed sets, where $i = \sigma(j'_\sigma)$, $B_1 \subseteq C \setminus M$, and $B_2 \subseteq M \setminus \{i\}$, $|B_2| = m-k$, and then summing over these sets, we obtain

$$\begin{aligned}\Pr(T_S = T_{k:M}) \;&=\; \sum_{B_1 \subseteq C \setminus M} \; \sum_{i \in M} \; \sum_{\substack{B_2 \subseteq M \setminus \{i\} \\ |B_2| = m-k}} \big(\phi(B_1 \cup B_2 \cup \{i\}) - \phi(B_1 \cup B_2)\big) \\ &\quad \times \sum_{\substack{\sigma \in \mathfrak{S}_n \,:\, \{\sigma(n-|B_1 \cup B_2|+1), \ldots, \sigma(n)\} = B_1 \cup B_2 \\ \sigma(n-|B_1 \cup B_2|) = i}} \Pr(T_{\sigma(1)} < \cdots < T_{\sigma(n)})\,.\end{aligned}$$

By (6) the inner sum is precisely $q_i(B_1 \cup B_2)$. We then conclude by setting $A = B_1 \cup B_2$. The second expression can be obtained by permuting the sums of the first expression. □

Example 3. Consider a 3-component system whose structure function is given by

$$\phi(x_1, x_2, x_3) \;=\; (x_1 \amalg x_2)\, x_3 \;=\; x_1 x_3 + x_2 x_3 - x_1 x_2 x_3\,.$$

For such a system, we have for instance

$$\begin{aligned} p_1^{\{1,3\}} \;&=\; \sum_{\substack{A \subseteq \{1,2,3\} \\ |\{1,3\} \setminus A| = 1}} \; \sum_{j \in \{1,3\} \setminus A} q_j(A)\, \Delta_j\, \phi(A) \\ &=\; q_1(\{3\}) + q_3(\{1\}) + q_3(\{1,2\}) \\ &=\; \Pr(T_2 < T_1 < T_3) + \Pr(T_2 < T_3 < T_1) + \Pr(T_3 < T_1 < T_2) + \Pr(T_3 < T_2 < T_1). \end{aligned}$$

Interestingly, Theorem 2 provides new expressions for p_k, namely

$$p_k \;=\; \sum_{\substack{A \subseteq C \\ |A| = n-k}} \; \sum_{j \in C \setminus A} q_j(A)\, \Delta_j\, \phi(A) \;=\; \sum_{j \in C} \; \sum_{\substack{A \subseteq C \setminus \{j\} \\ |A| = n-k}} q_j(A)\, \Delta_j\, \phi(A)\,.$$

Replacing $\Delta_j\, \phi(A)$ by $\phi(A \cup \{j\}) - \phi(A)$ in formula (7), we derive the following alternative expression for the probability p_k^M.

Corollary 4. *For every nonempty set $M \subseteq C$ and every $k \in \{1, \ldots, m\}$, we have*

$$p_k^M \;=\; \sum_{j \in M} \; \sum_{\substack{A \subseteq C \\ |(M \setminus A) \cup \{j\}| = k}} (-1)^{\{j\} \setminus A}\, q_j(A \setminus \{j\})\, \phi(A)\,. \tag{8}$$

Proof. The right-hand side of (8) can be written as

$$\begin{aligned} &\sum_{j \in M} \; \sum_{\substack{A \subseteq C,\ A \ni j \\ |M \setminus A| = k-1}} q_j(A \setminus \{j\})\, \phi(A) - \sum_{j \in M} \; \sum_{\substack{A \subseteq C,\ A \not\ni j \\ |M \setminus A| = k}} q_j(A)\, \phi(A) \\ &\quad =\; \sum_{j \in M} \; \sum_{\substack{A \subseteq C \setminus \{j\} \\ |M \setminus A| = k}} q_j(A)\, \phi(A \cup \{j\}) - \sum_{j \in M} \; \sum_{\substack{A \subseteq C \setminus \{j\} \\ |M \setminus A| = k}} q_j(A)\, \phi(A)\,, \end{aligned}$$

which is precisely the right-hand side of (7). □

We now provide an alternative linear expression for the probability p_k^M that generalizes formula (2). For every nonempty set $M\subseteq C$, define the set functions $q_M^{\downarrow}\colon 2^C\setminus\{\varnothing\}\to\mathbb{R}$ and $q_M^{\uparrow}\colon 2^C\setminus\{C\}\to\mathbb{R}$ by

$$q_M^{\downarrow}(A) ~=~ \sum_{j\in M\cap A} q_j(A\setminus\{j\}) ~=~ \Pr\Big(\exists\, j\in M : \max_{i\in C\setminus A} T_i < T_j = \min_{i\in A} T_i\Big)$$

and

$$q_M^{\uparrow}(A) ~=~ \sum_{j\in M\setminus A} q_j(A) ~=~ \Pr\Big(\exists\, j\in M : \max_{i\in C\setminus A} T_i = T_j < \min_{i\in A} T_j\Big),$$

respectively.

Corollary 5. *For every nonempty set $M\subseteq C$ and every $k\in\{1,\ldots,m\}$, we have*

$$p_k^M ~=~ \sum_{\substack{A\subseteq C\\ |M\cap A|=m-k+1}} q_M^{\downarrow}(A)\,\phi(A) - \sum_{\substack{A\subseteq C\\ |M\cap A|=m-k}} q_M^{\uparrow}(A)\,\phi(A)\,. \tag{9}$$

Proof. By (2) we have

$$\begin{aligned}
p_k^M &= \sum_{\substack{A\subseteq C\\ |M\setminus A|=k}} \sum_{j\in M\setminus A} q_j(A)\,\Delta_j\,\phi(A)\\
&= \sum_{\substack{A\subseteq C\\ |M\setminus A|=k}} \sum_{j\in M\setminus A} q_j(A)\,\phi(A\cup\{j\}) - \sum_{\substack{A\subseteq C\\ |M\setminus A|=k}} \sum_{j\in M\setminus A} q_j(A)\,\phi(A)\\
&= \sum_{\substack{A\subseteq C\\ |M\setminus A|=k-1}} \sum_{j\in M\cap A} q_j(A\setminus\{j\})\,\phi(A) - \sum_{\substack{A\subseteq C\\ |M\setminus A|=k}} q_M^{\uparrow}(A)\,\phi(A)\,,
\end{aligned}$$

which completes the proof. □

We end this section by providing an explicit linear expression for the probability p_k^M in terms of the Möbius transform of the structure function ϕ. Recall that the Möbius transform of ϕ is the set function $c_\phi\colon 2^C\to\mathbb{R}$ which gives the coefficients of the unique multilinear expression of the structure function, that is,

$$\phi(\mathbf{x}) ~=~ \sum_{A\subseteq C} c_\phi(A)\prod_{i\in A} x_i\,.$$

The conversion formulas between c_ϕ and ϕ are given by

$$c_\phi(A) ~=~ \sum_{B\subseteq A}(-1)^{|A|-|B|}\,\phi(B) \quad\text{and}\quad \phi(A) ~=~ \sum_{B\subseteq A} c_\phi(B)\,. \tag{10}$$

A very simple linear expression for p_k^M in terms of the Möbius transform c_ϕ is given in the following theorem.

Theorem 6. *For every nonempty set $M\subseteq C$ and every $k\in\{1,\ldots,m\}$, we have*

$$p_k^M ~=~ \sum_{\substack{A\subseteq C\\ |M\cap A|\leqslant m-k+1}} c_\phi(A)\,\Pr\Big(T_{k:M} = \min_{i\in A} T_i\Big),$$

or equivalently,

$$p_k^M ~=~ \sum_{A\subseteq C} c_\phi(A)\,\Pr\Big(T_{k:M} = \min_{i\in A} T_i\Big).$$

Proof. By substituting the second formula of (10) in (9) and then permuting the resulting sums, we obtain

$$\begin{aligned} p_k^M &= \sum_{\substack{A\subseteq C\\ |M\cap A|=m-k+1}} q_M^{\downarrow}(A)\sum_{B\subseteq A} m(B) - \sum_{\substack{A\subseteq C\\ |M\cap A|=s-k}} q_M^{\uparrow}(A)\sum_{B\subseteq A} m(B)\\ &= \sum_{\substack{B\subseteq C\\ |M\cap B|\leqslant m-k+1}} m(B)\sum_{\substack{A\supseteq B\\ |M\cap A|=m-k+1}} q_M^{\downarrow}(A) - \sum_{\substack{B\subseteq C\\ |M\cap B|\leqslant m-k}} m(B)\sum_{\substack{A\supseteq B\\ |M\cap A|=m-k}} q_M^{\uparrow}(A)\,. \end{aligned}$$

However, we have

$$\begin{aligned} \sum_{\substack{A\supseteq B\\ |M\cap A|=m-k+1}} q_M^{\downarrow}(A) &= \Pr\Big(\exists\, j\in M,\ \exists\, A\supseteq B,\ |M\cap A|=m-k+1 : \max_{i\in C\setminus A} T_i < T_j = \min_{i\in A} T_i\Big)\\ &= \Pr\Big(T_{k:M}\leqslant \min_{i\in B} T_i\Big) \end{aligned}$$

and

$$\begin{aligned} \sum_{\substack{A\supseteq B\\ |M\cap A|=m-k}} q_M^{\uparrow}(A) &= \Pr\Big(\exists\, j\in M,\ \exists\, A\supseteq B,\ |M\cap A|=m-k : \max_{i\in C\setminus A} T_i = T_j < \min_{i\in A} T_i\Big)\\ &= \Pr\Big(T_{k:M} < \min_{i\in B} T_i\Big)\,. \end{aligned}$$

Thus, we have

$$\begin{aligned} p_k^M &= \sum_{\substack{B\subseteq C\\ |M\cap B|\leqslant m-k+1}} m(B)\,\Pr\Big(T_{k:M}\leqslant \min_{i\in B} T_i\Big) - \sum_{\substack{B\subseteq C\\ |M\cap B|\leqslant m-k}} m(B)\,\Pr\Big(T_{k:M} < \min_{i\in B} T_i\Big)\\ &= \sum_{\substack{B\subseteq C\\ |M\cap B|= m-k+1}} m(B)\,\Pr\Big(T_{k:M}\leqslant \min_{i\in B} T_i\Big) + \sum_{\substack{B\subseteq C\\ |M\cap B|\leqslant m-k}} m(B)\,\Pr\Big(T_{k:M} = \min_{i\in B} T_i\Big)\,. \end{aligned}$$

We observe that we cannot have $T_{k:M} < \min_{i\in B} T_i$ if $|M\cap B| = m-k+1$. This proves the first formula of the theorem. To see that the second formula holds, just observe that we cannot have $T_{k:M} = \min_{i\in B} T_i$ if $|M\cap B| > m-k+1$. ☐

Example 7. Consider a system defined as in Example 3, that is, with a design given by the (multilinear) structure function

$$\phi(x_1,x_2,x_3) \;=\; x_1x_3 + x_2x_3 - x_1x_2x_3\,.$$

For such a system, we have for instance

$$\begin{aligned} p_1^{\{1,3\}} &= \sum_{A\subseteq\{1,2,3\}} c_{\phi}(A)\,\Pr\Big(T_{1:\{1,3\}} = \min_{i\in A} T_i\Big)\\ &= 1 + \Pr\Big(\min_{i\in\{1,3\}} T_i = \min_{i\in\{2,3\}} T_i\Big) - \Pr\Big(\min_{i\in\{1,3\}} T_i = \min_{i\in\{1,2,3\}} T_i\Big)\\ &= \Pr(T_2<T_1<T_3) + \Pr(T_2<T_3<T_1) + \Pr(T_3<T_1<T_2) + \Pr(T_3<T_2<T_1). \end{aligned}$$

Remark 2. We observe that the probability $\Pr(T_{k:M} = \min_{i\in A} T_i)$ is exactly the kth coordinate of the M-signature of the semicoherent system obtained from the current system by transforming the structure function into $\phi(\mathbf{x}) = \prod_{i\in A} x_i$. This result follows immediately from the fact that the modified system has lifetime $\min_{i\in A} T_i$.

From Theorem 6 we immediately derive the following corollary, which was already established in [6].

Corollary 8. *For every $k\in\{1,\ldots,n\}$ and every $j\in C$, we have*

$$p_k ~=~ \sum_{A\subseteq C} c_\phi(A)\,\Pr\Big(T_{k:n}=\min_{i\in A}T_i\Big)\quad\text{and}\quad I_{\mathrm{BP}}^{(j)} ~=~ \sum_{A\subseteq C} c_\phi(A)\,\Pr\Big(T_j=\min_{i\in A}T_i\Big).$$

3. Exchangeable component lifetimes

We now consider the special case when the functions q_j $(j\in C)$ satisfy the condition

$$q_j(A) ~=~ \frac{1}{n\binom{n-1}{|A|}}\,,\qquad A\subseteq C\smallsetminus\{j\}\,,\ j\in C\,. \tag{11}$$

It is easy to see that this condition holds whenever the lifetimes $T_1,\ldots,T_n$ are i.i.d. or, more generally, exchangeable (see [6]). In this case, for every nonempty subset $M\subseteq C$, we also have

$$q_M^{\downarrow}(A) ~=~ \frac{|M\cap A|}{n\binom{n-1}{|A|-1}}\,,\quad q_M^{\uparrow}(A) ~=~ \frac{|M\smallsetminus A|}{n\binom{n-1}{|A|}}\,,\quad\text{and}\quad q(A) ~=~ \frac{1}{\binom{n}{|A|}}\,.$$

Theorem 2 and Corollaries 4 and 5 are then immediately specialized to the following result. Here the probability p_k^M is naturally denoted by s_k^M.

Corollary 9. *Assume that the functions q_j $(j\in C)$ satisfy condition (11). For every nonempty set $M\subseteq C$ and every $k\in\{1,\ldots,m\}$, we have*

$$s_k^M ~=~ \sum_{\substack{A\subseteq C\\ |M\smallsetminus A|=k}}\frac{1}{n\binom{n-1}{|A|}}\sum_{j\in M\smallsetminus A}\Delta_j\,\phi(A) ~=~ \sum_{j\in M}\sum_{\substack{A\subseteq C\smallsetminus\{j\}\\ |M\smallsetminus A|=k}}\frac{1}{n\binom{n-1}{|A|}}\,\Delta_j\,\phi(A)\,, \tag{12}$$

$$s_k^M ~=~ \sum_{j\in M}\sum_{\substack{A\subseteq C\\ |(M\smallsetminus A)\cup\{j\}|=k}}(-1)^{\{j\}\smallsetminus A}\,\frac{1}{n\binom{n-1}{|A\smallsetminus\{j\}|}}\,\phi(A)\,,$$

and

$$s_k^M ~=~ \sum_{\substack{A\subseteq C\\ |M\cap A|=m-k+1}}\frac{m-k+1}{n\binom{n-1}{|A|-1}}\,\phi(A)-\sum_{\substack{A\subseteq C\\ |M\cap A|=m-k}}\frac{k}{n\binom{n-1}{|A|}}\,\phi(A)\,.$$

From (12) we immediately derive new expressions for s_k, namely

$$s_k ~=~ \sum_{\substack{A\subseteq C\\ |A|=n-k}}\frac{1}{n\binom{n-1}{|A|}}\sum_{j\in C\smallsetminus A}\Delta_j\,\phi(A) ~=~ \sum_{j\in C}\sum_{\substack{A\subseteq C\smallsetminus\{j\}\\ |A|=n-k}}\frac{1}{n\binom{n-1}{|A|}}\,\Delta_j\,\phi(A)\,.$$

An expression for s_k^M in terms of the Möbius transform of the structure function is given in the following result.

Corollary 10. *Assume that the functions q_j $(j\in C)$ satisfy condition (11). For every nonempty set $M\subseteq C$ and every $k\in\{1,\ldots,m\}$, we have*

$$s_k^M ~=~ \sum_{\substack{A\subseteq C\\ k-1\leqslant|M\smallsetminus A|\leqslant m-1}} c_\phi(A)\,\frac{m-|M\smallsetminus A|}{k}\,\frac{\binom{|M\smallsetminus A|}{k-1}}{\binom{|M\smallsetminus A|+|A|}{k}}\,,$$

or equivalently,

$$s_k^M ~=~ \sum_{A\subseteq C} c_\phi(A)\,\frac{m-|M\smallsetminus A|}{k}\,\frac{\binom{|M\smallsetminus A|}{k-1}}{\binom{|M\smallsetminus A|+|A|}{k}}\,.$$

Proof. For every $A \subseteq C$, let $\phi_A(\mathbf{x}) = \prod_{i\in A} x_i$. For every $B \subseteq C$, we then have $\Delta_j\, \phi_A(B) = 1$, if $j \in A$ and $A \setminus \{j\} \subseteq B$, and 0, otherwise.

Combining Remark 2 with (12), we then obtain

$$\begin{aligned}\Pr\Big(T_{k:M} = \min_{i\in A} T_i\Big) &= \sum_{j\in M}\ \sum_{\substack{B\subseteq C\setminus\{j\}\\ |M\setminus B|=k}} \frac{1}{n\binom{n-1}{|B|}}\,\Delta_j\,\phi_A(B)\\ &= \sum_{j\in M\cap A}\ \sum_{\substack{A\setminus\{j\}\subseteq B\subseteq C\setminus\{j\}\\ |M\setminus B|=k}} \frac{1}{n\binom{n-1}{|B|}}\end{aligned}$$

Partitioning A into $A_1 = A\cap M$ and $A_2 = A\setminus M$, the latter expression becomes

$$\begin{aligned}&\sum_{j\in A_1}\ \sum_{\substack{A_1\setminus\{j\}\subseteq B_1\subseteq M\setminus\{j\}\\ |B_1|=m-k}}\ \sum_{A_2\subseteq B_2\subseteq C\setminus M} \frac{1}{n\binom{n-1}{m-k+|B_2|}}\\ &= \sum_{j\in A_1}\binom{m-|A_1|}{m-k-|A_1|+1}\sum_{b_2=|A_2|}^{n-m}\binom{n-m-|A_2|}{b_2-|A_2|}\int_0^1 t^{m-k+b_2}\,(1-t)^{n-1-m+k-b_2}\,dt\\ &= |A_1|\binom{m-|A_1|}{k-1}\int_0^1 t^{m-k+|A_2|}\,(1-t)^{k-1}\,dt \;=\; \frac{|A_1|}{m+|A_2|}\,\frac{\binom{m-|A_1|}{k-1}}{\binom{m+|A_2|-1}{k-1}}\,.\end{aligned}$$

We then conclude by Theorem 6. □

From Corollary 10 we immediately derive the following result.

Corollary 11. *Assume that the functions q_j $(j \in C)$ satisfy condition (11). For every $k \in \{1,\ldots,n\}$ and every $j \in C$, we have*

$$s_k \;=\; \sum_{\substack{A\subseteq C\\ |A|\leqslant n-k+1}} c_\phi(A)\,\frac{|A|}{k}\,\frac{\binom{n-|A|}{k-1}}{\binom{n}{k}} \quad \text{and} \quad b_j \;=\; \sum_{A\subseteq C,\ A\ni j} c_\phi(A)\,\frac{1}{|A|}\,.$$

4. SUBSIGNATURES ASSOCIATED WITH MODULAR SETS

Suppose that the component lifetimes are i.i.d. and that the system contains a module (M,χ), where M is the corresponding modular set and $\chi\colon\{0,1\}^M \to \{0,1\}$ is the corresponding structure function. In this case the structure function of the system expresses through the composition

$$\phi(\mathbf{x}) \;=\; \psi\big(\chi(\mathbf{x}^M),\mathbf{x}^{C\setminus M}\big)\,, \tag{13}$$

where $\mathbf{x}^M = (x_i)_{i\in M}$, $\mathbf{x}^{C\setminus M} = (x_i)_{i\in C\setminus M}$, and, setting $C_M = (C\setminus M)\cup\{[M]\}$, the function $\psi\colon\{0,1\}^{C_M}\to\{0,1\}$ is the structure function of the reduced system (of $n-m+1$ components) obtained from the original system by considering the modular set M as a single macro-component $[M]$. For general background on modules, see [2, Chap. 1].

Denote by T_M the lifetime of the module. By (12) the signature of the module is an m-tuple whose kth coordinate is given by

$$\Pr(T_M = T_{k:M}) \;=\; \sum_{\substack{A\subseteq M\\ |A|=m-k}} \frac{1}{m\binom{m-1}{|A|}}\sum_{j\in M\setminus A}\Delta_j\,\chi(A)\,. \tag{14}$$

Denote by $h_\psi\colon [0,1]^{C_M}\to\mathbb{R}$ the reliability function of the structure function ψ, that is, the multilinear polynomial function

$$h_\psi(\mathbf{x}) ~=~ \sum_{A\subseteq C_M}\psi(A)\prod_{i\in A}x_i\prod_{i\in C_M\smallsetminus A}(1-x_i)\,.$$

We can easily see that the average value over $[0,1]$ of the function

$$(\partial_{[M]}h_\psi)(t,\ldots,t) ~=~ \sum_{A\subseteq C_M\smallsetminus\{[M]\}}t^{|A|}(1-t)^{n-m-|A|}\,\Delta_{[M]}\,\psi(A) \tag{15}$$

(see [8] for details) with respect to the uniform distribution is given by

$$\int_0^1(\partial_{[M]}h_\psi)(t,\ldots,t)\,dt ~=~ \sum_{A\subseteq C_M\smallsetminus\{[M]\}}\frac{1}{(n-m+1)\binom{n-m}{|A|}}\,\Delta_{[M]}\,\psi(A)\,.$$

By (3) this quantity is precisely the $[M]$-coordinate of the Barlow-Proschan importance index for the reduced system (C_M,ψ). Thus, we have

$$\int_0^1(\partial_{[M]}h_\psi)(t,\ldots,t)\,dt ~=~ \Pr(T_S=T_M)\,. \tag{16}$$

The following result shows that s_k^M factorizes into the product of $\Pr(T_M=T_{k:M})$ and the expected value of the function $(\partial_{[M]}h_\psi)(t,\ldots,t)$ with respect to a beta distribution. When M is a singleton, this result reduces to (16).

Theorem 12. *Assume that the components have i.i.d. lifetimes. For every nonempty modular set $M\subseteq C$ and every $k\in\{1,\ldots,m\}$, we have*

$$s_k^M ~=~ \Pr(T_M=T_{k:M})\int_0^1 r_{k,m}(t)\,(\partial_{[M]}h_\psi)(t,\ldots,t)\,dt\,,$$

where ψ is the structure function defined in (13) and $r_{k,m}(t)$ is the p.d.f. of the beta distribution on $[0,1]$ with parameters $\alpha=m-k+1$ and $\beta=k$.

Proof. Let $h_\phi\colon[0,1]^C\to\mathbb{R}$ and $h_\chi\colon[0,1]^M\to\mathbb{R}$ be the reliability functions of ϕ and χ, respectively. We then have

$$h_\phi(\mathbf{x}) ~=~ h_\psi\big(h_\chi(\mathbf{x}^M),\mathbf{x}^{C\smallsetminus M}\big)\,.$$

By the chain rule it follows that, for every $j\in M$,

$$\partial_j h_\phi(\mathbf{x}) ~=~ (\partial_{[M]}h_\psi)\big(h_\chi(\mathbf{x}^M),\mathbf{x}^{C\smallsetminus M}\big)\,\partial_j h_\chi(\mathbf{x}^M)\,.$$

Since $\partial_{[M]}h_\psi$ does not depend on its $[M]$-variable, we have

$$(\partial_{[M]}h_\psi)\big(h_\chi(\mathbf{x}^M),\mathbf{x}^{C\smallsetminus M}\big) ~=~ (\partial_{[M]}h_\psi)\big(0,\mathbf{x}^{C\smallsetminus M}\big)\,.$$

Since the reliability functions are multilinear, for every $A_1\subseteq M$ and every $A_2\subseteq C\smallsetminus M$ we immediately obtain the factorization

$$\Delta_j\,\phi(A_1\cup A_2) ~=~ \Delta_{[M]}\,\psi(A_2)\,\Delta_j\,\chi(A_1).$$

Therefore, by (12) and (14) we obtain

$$\begin{aligned} s_k^M &= \sum_{\substack{A_1\subseteq M\\ |A_1|=m-k}} \sum_{A_2\subseteq C\smallsetminus M} \frac{1}{n\binom{n-1}{m-k+|A_2|}} \sum_{j\in M\smallsetminus A_1} \Delta_j\,\phi(A_1\cup A_2)\\ &= \Bigg(\sum_{\substack{A_1\subseteq M\\ |A_1|=m-k}} \frac{1}{m\binom{m-1}{m-k}} \sum_{j\in M\smallsetminus A_1} \Delta_j\,\chi(A_1)\Bigg)\Bigg(\sum_{A_2\subseteq C\smallsetminus M} \frac{m\binom{m-1}{m-k}}{n\binom{n-1}{m-k+|A_2|}}\,\Delta_{[M]}\,\psi(A_2)\Bigg)\\ &= \Pr(T_M=T_{k:M}) \sum_{A_2\subseteq C\smallsetminus M} \frac{\int_0^1 t^{m-k+|A_2|}\,(1-t)^{k-1+n-m-|A_2|}\,dt}{\int_0^1 t^{m-k}\,(1-t)^{k-1}\,dt}\,\Delta_{[M]}\,\psi(A_2) \end{aligned}$$

and we conclude by (15). □

Interestingly, Theorem 12 shows that in the i.i.d. case the ratio $s_k^M/\Pr(T_M = T_{k:M})$ is independent of the structure χ of the module (assuming $\Pr(T_M = T_{k:M}) \neq 0$). Thus, if the components of the module are reorganized so that χ is modified into χ' and if the probability $\Pr(T_M = T_{k:M})$ remains unchanged, then so does s_k^M.

ACKNOWLEDGMENTS

The author wish to thank Pierre Mathonet for fruitful discussions. This research is supported by the internal research project F1R-MTH-PUL-12RDO2 of the University of Luxembourg.

REFERENCES

[1] R.E. Barlow and F. Proschan. Importance of system components and fault tree events. *Stochastic Processes and Their Applications*, 3:153–172, 1975.

[2] R.E. Barlow and F. Proschan. *Statistical theory of reliability and life testing.* To Begin With, Silver Spring, MD, 1981.

[3] P. J. Boland. Signatures of indirect majority systems. *J. Appl. Prob.*, 38:597–603, 2001.

[4] S. Iyer. The Barlow-Proschan importance and its generalizations with dependent components. *Stochastic Process. Appl.*, 42:353–359, 1992.

[5] J.-L. Marichal and P. Mathonet. Extensions of system signatures to dependent lifetimes: Explicit expressions and interpretations. *J. Multivariate Analysis*, 102(5):931–936, 2011.

[6] J.-L. Marichal and P. Mathonet. On the extensions of Barlow-Proschan importance index and system signature to dependent lifetimes. *J. of Multivariate Analysis*, 115:48–56, 2013.

[7] J.-L. Marichal, P. Mathonet, and T. Waldhauser. On signature-based expressions of system reliability. *J. Multivariate Analysis*, 102(10): 1410–1416, 2011.

[8] G. Owen. Multilinear extensions of games. *Management Sci.*, 18:P64–P79, 1972.

[9] F.J. Samaniego. On closure of the IFR class under formation of coherent systems. *IEEE Trans. Reliability Theory*, 34:69–72, 1985.

[10] F.J. Samaniego. *System signatures and their applications in engineering reliability.* Int. Series in Operations Research & Management Science, 110. New York: Springer, 2007.

MATHEMATICS RESEARCH UNIT, FSTC, UNIVERSITY OF LUXEMBOURG, 6, RUE COUDENHOVE-KALERGI, L-1359 LUXEMBOURG, LUXEMBOURG

E-mail address: `jean-luc.marichal[at]uni.lu`